

What is claimed is:

1. A method for increasing the hybridization rate of nucleic acids in a nucleic acid assay, said method comprising:
 - a) attaching probe nucleic acid molecules of known sequence to a solid support;
 - b) labeling nucleic acid target molecules with paramagnetic labels;
 - c) contacting the labeled target molecules with the solid support;
 - d) activating a magnetic field whereby the labeled molecules are attracted to the solid support;
 - e) washing the support and inverting the polarity of the magnetic field to remove any unbound or nonspecifically bound molecules; and
 - f) detecting the hybridized target nucleic acid molecules.
2. A method of claim 1 in which the solid support is selected from the group consisting of silicon, glass, and metals.
3. A method of claim 2 in which the solid support is or is coated with a metal selected from the group consisting of silver, copper, gold, platinum (II), mercury, mercury (II), thallium, cadmium (II), platinum (IV) and palladium (II).
4. A method of claim 1 in which the paramagnetic labels comprise superparamagnetic particles, having a diameter of from about 1 to about 10 nanometers.
5. A method of claim 1 in which the paramagnetic labels comprise paramagnetic porphyrins.

6 ~~5.~~ A method of claim 1 in which the paramagnetic labels are attached to the nucleic acid molecules using cleavable conjugating molecules.

7 ~~6.~~
5 A method of claim 1 in which the nucleic acid molecules are oligonucleotides, genomic DNA, cDNA, RNA or fragments thereof.

8 ~~7.~~ A method of claim 1 in which at least one member of a complementary pair is labeled with a fluorescent detection molecule.

10 ~~8.~~ A method for increasing the hybridization rate of nucleic acids in a nucleic acid assay, said method comprising:

- US 5,927,425 A 11/19001
15
20
- a) attaching nucleic acid target molecules to a solid support;
 - b) labeling nucleic acid molecules of known sequence with paramagnetic labels;
 - c) contacting the support with the labeled nucleic acid molecules of known sequence;
 - d) activating a magnetic field whereby the labeled molecules are attracted to the solid support;
 - e) washing the support and inverting the polarity of the magnetic field to remove any unbound or nonspecifically bound molecules; and
 - f) detecting the hybridized target nucleic acid molecules.

10 ~~9.~~ A method of claim 9 in which the solid support is selected from the group consisting of silicon, glass, and metals.

25 ~~10.~~
~~11.~~
16 A method of claim 10 in which the solid support is or is coated with a metal selected from the group consisting of silver, copper, gold, platinum (II), mercury, mercury (II), thallium, cadmium (II), platinum (IV) and palladium (II).

12. A method of claim 9 in which the paramagnetic labels comprise superparamagnetic particles, having a diameter of from about 1 to about 10 nanometers.

13. A method of claim 9 in which the paramagnetic labels comprise paramagnetic porphyrins.

14. A method of claim 9 in which the paramagnetic labels are attached to the nucleic acid molecules using cleavable conjugating molecules.

15. A method of claim 9 in which the nucleic acid molecules are oligonucleotides, genomic DNA, cDNA, RNA or fragments thereof.

16. A method of claim 9 in which at least one member of a complementary pair is labeled with a fluorescent detection molecule.